

INVENTORS: Max F. Rothschild  
Kwan Suk Kim  
Rebecca S. Emnett

5 TITLE: GENETIC MARKERS FOR IMPROVED MEAT CHARACTERISTICS  
IN ANIMALS

GRANT REFERENCE CLAUSE

10 This invention was supported at least in part by grants  
from the United States Department of Agriculture through the  
Iowa Agriculture and Home Economics Experiment Station  
(IAHEES) and Project Number IOW03148 (Hatch Funds). The  
United States government may have certain rights in this  
invention.

15

FIELD OF THE INVENTION

The present invention relates to a method of genetically  
evaluating animals by assaying for the presence of at least  
one genetic marker which is indicative of one or more traits  
20 associated with meat quality. In particular, the method  
analyzes for variation in the melanocortin-4 receptor (MC4R)  
gene or other variations associated therewith which are  
indicative of these favorable traits.

25 BACKGROUND OF THE INVENTION

Genetic differences exist among individual animals as  
well as among breeds which can be exploited by breeding  
techniques to achieve animals with desirable characteristics.  
For example, Chinese breeds are known for reaching puberty at  
30 an early age and for their large litter size, while American  
breeds are known for their greater growth rates and leanness.  
Often, however, heritability for desired traits is low, and  
standard breeding methods which select individuals based upon  
phenotypic variations do not take fully into account genetic  
35 variability or complex gene interactions which exist.

Restriction fragment length polymorphism (RFLP) analysis  
has been used by several groups to study pig DNA. Jung et

al., Theor. Appl. Genet., 77:271-274 (1989), incorporated herein by reference, discloses the use of RFLP techniques to show genetic variability between two pig breeds.

Polymorphism was demonstrated for swine leukocyte antigen

5 (SLA) Class I genes in these breeds. Hoganson et al.,  
Abstract for Annual Meeting of Midwestern Section of the  
American Society of Animal Science, March 26-28, 1990,  
incorporated herein by reference, reports on the polymorphism  
of swine major histocompatibility complex (MHC) genes for  
10 Chinese pigs, also demonstrated by RFLP analysis. Jung et  
al., Theor. Appl. Genet., 77:271-274 (1989), incorporated  
herein by reference, reports on RFLP analysis of SLA Class I  
genes in certain boars. The authors state that the results  
suggest that there may be an association between swine  
15 SLA/MHC Class I genes and production and performance traits.  
They further state that the use of SLA Class I restriction  
fragments, as genetic markers, may have potential in the  
future for improving pig growth performance.

The ability to follow a specific favorable genetic  
20 allele involves a novel and lengthy process of the  
identification of a DNA molecular marker for a major effect  
gene. The marker may be linked to a single gene with a major  
effect or linked to a number of genes with additive effects.  
DNA markers have several advantages; segregation is easy to  
25 measure and is unambiguous, and DNA markers are co-dominant,  
i.e., heterozygous and homozygous animals can be  
distinctively identified. Once a marker system is  
established selection decisions could be made very easily,  
since DNA markers can be assayed any time after a tissue or  
30 blood sample can be collected from the individual infant  
animal, or even an embryo.

The use of genetic differences in receptor genes has  
become a valuable marker system for selection. For example,  
United States Patents 5,550,024 and 5,374,526 issued to  
35 Rothschild et al. disclose a polymorphism in the pig estrogen  
receptor gene which is associated with larger litter size,  
the disclosure of which is incorporated herein by reference.

United States number 5,935,784 discloses polymorphic markers in the pig prolactin receptor gene which are associated with larger litter size and overall reproductive efficiency. Perhaps one of the most important characteristics for any meat producing animal is meat quality. Meat quality is a difficult characteristic to assess, as many different aspects, both objective and subjective, make up the overall trait. The list of factors which determine quality in meat, as with other foods, is rather long (Wood et al., Proceedings of The Nutrition Society (1999) 58:363-70). It includes freedom from microbiological hazards (food safety) and prevention of animal exploitation (animal welfare). It also includes the sensory appeal of meat, i.e. its taste or eating quality, and perceived healthiness, especially in relation to the amount and type of fat.

The quality of raw pig meat is influenced by a large number of genetic and non-genetic factors. The latter include farm, transport, slaughter and processing conditions. Meat scientists have performed a substantial amount of research on these factors, which has led to considerable quality improvement. Part of the research has also been dedicated to the genetic background of the pigs, and several studies have revealed the importance of genetic factors. This has made the industry aware that selective breeding of pigs and the use of gene technology can play an important role in enhancing pork quality.

Information at DNA level can help to fix a specific major gene, but it can also assist the selection of quantitative trait for which we already select. Molecular information in addition to phenotypic data can increase the accuracy of selection and therefore the selection response. The size of the extra response in such a Marker Assisted Selection (MAS) program has been considered by many workers from a theoretical point of view. In general terms, MAS is more beneficial for traits with a low heritability and which are expensive to measure phenotypically. Meat quality in particular qualifies as an excellent opportunity to utilize

MAS. For example, Meuwissen, T.H.E. and Goddard, M.E. (1996) "The use of Marker Haplotypes in Animal Breeding Schemes", Genet. Sel. Evol., 28 161-176 considered the impact of Marker Assisted Selection for traits such as reproduction and meat  
5 quality that are difficult to progress using traditional methods. their results are extremely encouraging, showing that for traits such as meat quality, where the trait is measured after slaughter, an additional response of up to 64% could be achieved.

10 Indeed, the best approach to genetically improve meat quality is to find relevant DNA-markers directly in the population under selection. Meat quality measurements can be performed continuously on some animals from the nucleus populations of breeding organizations. Since a full  
15 assessment of meat quality can only be done after slaughter, the data must be collected on culled animals and cannot be obtained on potential breeding animals.

This phenotypic meat quality data is collected in order to enable the detection of relevant DNA markers, and to  
20 validate markers from experimental populations or to test candidate genes. Significant markers or genes can then be included directly in the selection process. An advantage of the molecular information is that we can obtain it already at very young age of the breeding animal, which means that  
25 animals can be preselected based on DNA markers before the growing performance test is completed. This is a great advantage for the overall testing and selection system.

It can be seen from the foregoing that a need exists for a method for improving meat quality characteristics in  
30 animals by identifying and selecting animals with the improved meat characteristics.

An object of the present invention is to provide a genetic marker based on or within the MC4R gene which is indicative of favorable meat characteristics such as those  
35 evidenced by pH, marbling, color and drip loss.

Another object of the invention is to provide an assay for determining the presence of this genetic marker.

A further object of the invention is to provide a method of evaluating animals that increases accuracy of selection and breeding methods for the desired traits.

Yet another object of the invention is to provide a PCR  
5 amplification test which will greatly expedite the determination of presence of the marker.

An additional object of the invention is to provide a kit for evaluating a sample of animal DNA for the identified genetic marker.

10 These and other objects, features, and advantages will become apparent after review of the following description and claims of the invention which follow.

#### SUMMARY OF THE INVENTION

15 This invention relates to the discovery of a polymorphism within the melanocortin-4 receptor (MC4R) gene which is associated with meat quality traits in animals. This gene is highly conserved among species and it is expected that the different alleles disclosed herein will  
20 also correlate with variability in this gene in other meat producing animals such as bovine, sheep, chicken, etc. This polymorphic site has been previously described in an earlier patent application PCT/US99/16862, publication number WO 00/06777 the disclosure of which is incorporated herein. In  
25 the earlier application this site was found to significantly correlate with weight gain and feed intake, in other words, traits involving growth rate of the pig. Surprisingly, as fast growth is generally considered to be negatively correlated with meat quality, the marker has now been shown  
30 to correlate with favorable meat characteristics such as pH level, marbling, color, and drip loss. These multigenic characteristics have been previously difficult to associate with quantitative trait loci and current improvements in meat characteristics have centered around understanding and  
35 controlling the numerous factors i.e. on a farm, transport, and/or slaughter plant handling influencing meat quality including the incidence of PSE (pale, soft, exudative), RSE

(red, soft, exudative), and DFD (dark, firm, dry) meat.

According to the invention, the association of the MC4R polymorphism with the these trait(s) enables genetic markers to be identified for specific breeds or genetic lines to

5 identify animals with favorable meat characteristics early in the animal's life.

The marker genotype consists of a polymorphism within the MC4R gene that results in a guanine to adenine transition and a missense mutation of aspartic acid(D) codon (GAU) into  
10 asparagine(N) codon (AAU) at a position corresponding to amino acid position 298 of the human MC4R protein resulting in a *TaqI* restriction site in one allele of the gene. In one embodiment of the invention a *TaqI* restriction pattern which identifies the polymorphism is used to assay for the presence  
15 or absence of markers associated with the desirable meat traits. The invention includes assays for detection of the marker, or markers linked thereto as well as the sequence characterization of the polymorphism and includes novel sequences in the MC4R gene which may be used to design  
20 amplification primers for such an assay (SEQ ID NO:1). Additionally, the invention includes a method for using the assay in breeding programs for animal selection and a kit for performing the assay.

#### 25 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is the sequence listing for MC4R in pigs (SEQ ID NO:1). "X" represents the site of the polymorphism.

Figure 2 represents a comparison of the DNA sequence between the human (SEQ ID NO:2) and the porcine (SEQ ID NO:3)  
30 MC4R gene.

Figure 3 represents a comparison of the amino acid sequence between the human (SEQ ID NO:4) and the porcine (SEQ ID NO:5) MC4R gene.

Figures 4a, 4b, and 4c are linkage reports for MC4R from  
35 CRI-MAP.

Figure 5 depicts partial nucleotide and amino acid sequences (SEQ ID NO:12) of the porcine MC4R gene. The amino

acid translation shows an amino acid substitution at codon 298.

Figure 6 depicts multiple-alignments of the putative seventh transmembrane domain of porcine MC4R with other MCRs and GPCRs. The "\*" represents the predicted sequence positions for porcine MC4R. The other amino acid sequences were obtained from the GenBank database (accession numbers P32245, P70596, P41983, P56451, P34974, P41968, P33033, Q01718, Q01726, Q28031, AF011466, P21554, P18089, P30680, P47211). The missense variant in porcine MC4R substituted amino acid N for D in the position marked with an arrow. The Asp (D) residue is highly conserved among MCRs, and the Asn (N) residue is well conserved in most other GPCRs.

#### DETAILED DESCRIPTION OF THE INVENTION

The melanocortin-4 receptor (MC4R) has been shown to be an important mediator of long term weight homeostasis. MC4R antagonists can increase food intake and body weight during chronic administration. Skuladottir, G.V., et al., "Long term orexigenic effect of a novel melanocortin 4 receptor selective antagonist", British J. of Pharm., 126(1):27-34 (1999).

Lu et al., Nature (Oct 27, 1994), 371 (6500):799-802 suggested that the melanocortin receptors are involved in controlling food intake and energy balance through studying its antagonism to the *agouti* obesity syndrome. Huszar et al., Cell (Jan. 10, 1997) 88(1):131-41 found that inactivation of the melanocortin-4 receptor gene (MC4R) resulted in a maturity onset obesity syndrome in mice and demonstrated a major role of MC4R protein in the regulation of energy balance related to the *agouti* obesity syndrome. In addition, the MC4R protein mediates the effects of leptin, one of the important signaling molecules in energy homeostasis (Seeley et al. 1997).

According to the present invention, a variant or polymorphism in the MC4R gene has been located, and this genetic variability is associated with phenotypic differences

in the porcine meat quality traits as evidenced by pH, marbling, color and drip loss.

In one embodiment of the invention, an assay is provided for detection of presence of a desirable genotype in animals.

5 The assay involves assaying the genomic DNA purified from blood, tissue, semen, or other convenient source of genetic material by the use of primers and standard techniques, such as the polymerase chain reaction (PCR), then digesting the DNA with a restriction enzyme (e.g., *Taq I* or other enzyme  
10 which cleaves at the same G → A site) so as to yield gene fragments of varying lengths, and separating at least some of the fragments from others (e.g., using electrophoresis).

The fragments may also be detected by hybridizing with a nucleotide probe (e.g., radio-labeled cDNA probes) that  
15 contains all or at least a portion of the MC4R gene cDNA sequence to the separated fragments and comparing the results of the hybridization with assay results for a gene sequence known to have the marker or a sequence known to not have the marker. Selection and use of probes for detection of MC4R  
20 sequences based on the known and disclosed MC4R sequences is generally known to those skilled in the art. The probe may be any sequence which will hybridize to the separated digestion products and allow for detection.

Another embodiment of the invention provides a kit for  
25 assaying the presence in a MC4R gene sequence of a genetic marker. The marker being indicative of heritable traits of meat quality characteristics. The kit in a preferred embodiment also includes novel PCR primers comprising 4-30 contiguous bases on either side of the polymorphism to  
30 provide an amplification system allowing for detection of the G → A Transition polymorphism by PCR digestion of PCR products. The sequence surrounding the polymorphic site is shown in SEQ ID NO:1, Figure 1. Several primers have also been disclosed including SEQ ID NOS:6 and 7, SEQ ID NOS:10  
35 and 11; and mapping primers 8 and 9. The preferred primers are SEQ ID NO:10 and SEQ ID NO:11.



A further embodiment comprises a breeding method whereby an assay of the above type is conducted on a plurality of DNA samples from different animals or animal embryos to be selected from and based on the results, certain animals are  
5 either selected or dropped out of the breeding program.

According to the invention, in a preferred embodiment, the polymorphism in the MC4R gene identifiable by the *Taq I* restriction pattern, is disclosed. As is known in the art, restriction patterns are not exact determinants of the size  
10 of fragments and are only approximate. When the primers SEQ ID NOS:6 and 7 are used the polymorphism is identifiable by three bands from a *Taq I* digestion of the PCR product, 466, 225, and 76 base pairs (bp) for one homozygous genotype (allele 1); two bands, 542 and 225 bp for another homozygous  
15 genotype (allele 2); and four bands for the heterozygous genotype (542, 466, 225, and 76 bp). When the preferred primers are used, SEQ ID NOS:10 and 11 the bands upon *taq* digestion include 156 and 70 bp for allele 1 and one 226bp fragment for allele 2. Those of skill in the art will  
20 appreciate that the design of alternate primers PCR conditions and restriction patterns for identifying the presence of allele 2 using the MC4R sequence data herein or other data for closely linked loci represent nothing more than routine optimization of parameters and are intended to  
25 be within the scope of the invention. The marker for improved meat characteristics as evidenced by all four meat quality measurements observed herein (allele 2). The allele 2 genotype was previously associated with faster growth rate. This is surprising because the current state of the art  
30 concluded that there is a negative correlation between growth rate and meat quality.

In addition, the polymorphism associated with the pattern has been identified at the nucleotide level. The polymorphic *Taq I* site was sequenced along with the general  
35 surrounding area. See SEQ ID NO: 1. The sequences surrounding the polymorphism have facilitated the development of a PCR test in which a primer of about 4-30 contiguous

bases taken from the sequence immediately adjacent to the polymorphism is used in connection with a polymerase chain reaction to greatly amplify the region before treatment with the *Taq I* restriction enzyme. The primers need not be the exact complement; substantially equivalent sequences are acceptable.

From sequence data, it was observed that in allele 2 a guanine is substituted with an adenine at position 678 of the PCR product shown in Figure 1 or at position 298 of the analogous human MC4R amino acid of the MC4R protein changing the aspartic acid codon (GAU) into an asparagine codon (AAU). The PCR test for the polymorphism used a forward primer of 5'-TGG CAA TAG CCA AGA ACA AG-3' (SEQ. ID NO: 6) and a reverse primer of 5'-CAG GGG ATA GCA ACA GAT GA-3' (SEQ. ID NO: 7). Pig specific primers used for physical mapping were a forward primer of 5'-TTA AGT GGA GGA AGA AGG-3' (SEQ. ID NO: 8) and a reverse primer of 5'-CAT TAT GAC AGT TAA GCG G-3' (SEQ ID NO:9). The resulting amplified product of about 750 bp, when digested with *Taq I*, results in allelic fragments of 466, 225, and 76 bp (allele 1) or 542 and 225 bp (allele 2). The most preferred primers resulting in either 2 or 1 fragment after *Taq I* digestion are SEQ ID NOS:10 and 11. Allele 1 generates fragments of 156 and 70 base pairs while allele 2 generates a single 226 bp fragment.

The marker may be identified by any method known to one of ordinary skill in the art which identifies the presence or absence of the particular allele or marker, including for example, single-strand conformation polymorphism analysis (SSCP), RFLP analysis, heteroduplex analysis, denaturing gradient gel electrophoresis, allelic PCR, temperature gradient electrophoresis, ligase chain reaction, direct sequencing, minisequencing, nucleic acid hybridization, and micro-array-type detection of the MC4R gene and examination for the polymorphic site. Yet another technique includes an Invader Assay which includes isothermal amplification that relies on a catalytic release of fluorescence. See Third

Wave Technology at [www.twt.com](http://www.twt.com) all of which are intended to be within the scope of the invention.

One or more additional restriction enzymes and/or probes and/or primers can be used. Additional enzymes, constructed  
5 probes, and primers can be determined by routine experimentation by those of ordinary skill in the art and are intended to be within the scope of the invention.

Other possible techniques include non-gel systems such as TaqMan™ (Perkin Elmer). In this system, oligonucleotide  
10 PCR primers are designed that flank the mutation in question and allow PCR amplification of the region. A third oligonucleotide probe is then designed to hybridize to the region containing the base subject to change between different alleles of the gene. This probe is labeled with  
15 fluorescent dyes at both the 5' and 3' ends. These dyes are chosen such that while in this proximity to each other the fluorescence of one of them is quenched by the other and cannot be detected. Extension by Taq DNA polymerase from the PCR primer positioned 5' on the template relative to the  
20 probe leads to the cleavage of the dye attached to the 5' end of the annealed probe through the 5' nuclease activity of the Taq DNA polymerase. This removes the quenching effect allowing detection of the fluorescence from the dye at the 3' end of the probe. The discrimination between different DNA  
25 sequences arises through the fact that if the hybridization of the probe to the template molecule is not complete, i.e., there is a mismatch of some form, the cleavage of the dye does not take place. Thus, only if the nucleotide sequence of the oligonucleotide probe is completely complementary to  
30 the template molecule to which it is bound will quenching be removed. A reaction mix can contain two different probe sequences each designed against different alleles that might be present, thus, allowing the detection of both alleles in one reaction.

35 Though the use of RFLPs is one method of detecting the polymorphism, other methods known to one of ordinary skill in the art may be used. Such methods include ones that analyze

the polymorphic gene product and detect polymorphisms by detecting the resulting differences in the gene product.

Though the preferred method of separating restriction fragments is gel electrophoresis, other alternative methods  
5 known to one skilled in the art may be used to separate and determine the size of the restriction fragments.

It is possible to indirectly select for the polymorphism with alternative DNA markers and these methods are also within the scope of the invention. One can establish a  
10 linkage between specific alleles of alternative DNA markers and alleles of DNA markers known to be associated with the MC4R gene which have previously been shown to be associated with a particular trait. Examples of markers on the published PiGMap chromosome map which are linked to the MC4R  
15 gene include S0331, BHT0433, and S0313. This is also true for other species as well, for example in human the MC4R gene is located at chromosome 18q21.3-q22.

The reagents suitable for applying the methods of the present invention may be packaged into convenient kits. The  
20 kits provide the necessary materials, packaged into suitable containers. At a minimum, the kit contains a reagent that identifies the polymorphism in the MC4R gene that is associated with the trait of meat quality. Preferably, the reagent that identifies the polymorphism is a PCR set (a set  
25 of primers, DNA polymerase, and four nucleoside triphosphates) that hybridize with the MC4R gene or a fragment thereof. Preferably, the PCR set and restriction enzyme that cleaves the MC4R sequence in at least one place are included in the kit. Preferably, the kit further  
30 comprises additional means, such as buffers or reagents, for detecting or measuring the detectable entity or providing a control. Other reagents used for hybridization, prehybridization, DNA extraction, visualization, and similar purposes may also be included, if desired.

35 The genetic markers, methods, and kits of the invention are useful in a breeding program to identify and/or to select for meat characteristics in a breed, line, or population of

animals. Continuous selection and breeding of animals that are at least heterozygous and preferably homozygous for the desired polymorphism associated with the particular trait would lead to a breed, line, or population having those  
5 desired traits. Thus, the marker is a selection tool.

The following examples are offered to illustrate, but not limit the invention.

#### EXAMPLE 1

Melanocortin 4 Receptor PCR-RFLP Test - *TaqI* polymorphism and  
Genetic Linkage Mapping of MC4R Gene

##### Primers:

15 Primers were designed from homologous regions of human and rat MC4R sequences (Genbank Accession No. s77415 and u67863, respectively). These primers were used to amplify a 750-bp sequence of the porcine MC4R gene.

20 MC4R1: 5' TGG CAA TAG CCA AGA ACA AG 3' (SEQ ID NO:6)  
MC4R4: 5' CAG GGG ATA GCA ACA GAT GA 3' (SEQ ID NO:7)

##### PCR Conditions:

Mix 1:	10X Promega Buffer	1.0 $\mu$ L
	25 mM MgCl <sub>2</sub>	0.6 $\mu$ L
	dNTPs mix (2.5mM each)	0.5 $\mu$ L
	25 pmol/ $\mu$ L MC4R1	0.1 $\mu$ L
	25 pmol/ $\mu$ L MC4R4	0.1 $\mu$ L
	dd sterile H <sub>2</sub> O	7.5 $\mu$ L
	<i>Taq</i> Polymerase (5 U/ $\mu$ L)	0.07 $\mu$ L
	Genomic DNA (12.5 ng/ $\mu$ L)	1.0 $\mu$ L

25 Ten  $\mu$ L of Mix 1 and DNA were combined in reaction tube, then overlaid with mineral oil. The following PCR program was run: 94°C for 2 min.; 35 cycles of 94°C for 30 sec.; 58°C 1 min., and 72°C 1 min. 30 sec.; followed by a final extension at 72°C for 15 min.

Five  $\mu$ l of the PCR reaction product was checked on a standard 1% agarose gel to confirm amplification success and clean negative control. Product size is approximately 750 base pairs. Digestion was performed by the following

5 procedure.

<u>TaqI Digestion Reaction</u>	<u>10 <math>\mu</math>L reaction</u>
PCR product	5.0 $\mu$ L
10X TaqI NE Buffer	1.0 $\mu$ L
BSA (10mg/ml)	0.1 $\mu$ L
TaqI enzyme (20 U/ $\mu$ L)	0.5 $\mu$ L
dd sterile H <sub>2</sub> O	3.4 $\mu$ L

A cocktail of the buffer, enzyme, BSA, and water was made. Five  $\mu$ L was added to each reaction tube containing the  
10 DNA. The mixture was then incubated at 65°C for at least 4 hours to overnight. Loading dye was mixed with the digestion reaction and the total volume was loaded on a 3% agarose gel. The major bands for allele 1 are about 466, 225, and 76 bp. The allele 2 genotype bands are 542 and 225 bp. The  
15 heterozygote genotype has both allele 1 and allele 2.

### Results

The amplified PCR product is about 750 bp. The sequence of the PCR product confirmed that the PCR product is MC4R  
20 gene with 97.6%, and 92.2% identities at the amino acid and DNA level, respectively, to corresponding human sequences. (see Figs. 2 and 3).

The TaqI digestion of the PCR product produced allelic fragments of 466, 225, and 76 bp (allele 1), or 542 and 225  
25 bp (allele 2). The heterozygote genotype has both types of alleles. Mendelian inheritance was observed in three three-generation international reference families, which were used to map this gene by linkage analysis.

The polymorphism between allele 1 and allele 2 resulting  
30 from a G  $\rightarrow$  A transition at position 678 of the PCR product revealed a missense mutation of Aspartic acid codon (GAU)

into Asparagine codon (AAU) at position 298 amino acid of MC4R protein. (See Figure 1, SEQ ID NO:1).

Allele frequencies were determined by genotyping of DNA samples from a small number of animals from different breeds (Table 1). Allele 1 was observed with a frequency of 1 in Meishan, but was not observed or observed at very low frequency in Hampshire and Yorkshire. The frequencies of allele 1 in Landrace and Chester White were 0.5, respectively.

Figures 2 and 3 illustrate the differences between the DNA and amino acid sequences of the human and porcine MC4R gene (SEQ ID NOS:2-5).

TABLE 1  
The Frequency of Allele 1 in Different Pig Breeds

Breed	# Animals	Freq. Allele 1
Meishan	8	1
Large White	8	0.56
Yorkshire	6	0.08
Hampshire	5	0
Landrace	4	0.5
Chester White	4	0.5
Minzu	2	1
Wild Boar	2	1

#### Linkage Analyses

Two-point and multi-point linkage analyses were performed on the genotypes of international reference families. See Figs. 4a-4c. The data were analyzed by using the CRI-MAP program. MC4R was significantly linked to several markers on porcine chromosome (SSC) 1. The most closely linked markers (recombination fraction and LOD score in parentheses) are SO331 (0.02, 21.97), BHT0433 (0.02, 21.32), and SO313 (0.00, 17.76) by two-point linkage analysis. A multi-point linkage analysis produced the best

map order of markers and MC4R (with distance in Kosambi cM):  
KGF-5.8-CAPN3-2.5-MEF2A-6.1-MC4R-5.6-SO313.

Somatic cell hybrid panel of pig and rodent was used to  
assign MC4R to a cytogenetic region. PCR products from pig  
specific primers were amplified in clones 7, 8, 16, 18, and  
19. MC4R was localized to SSC1q 22-27.

## EXAMPLE 2

### Discovery of a Missense Variant of the Porcine Melanocortin-4 Receptor (MC4R) Gene

To determine if there was an association of this MC4R  
polymorphism with phenotypic variation the mutation was  
tested in a large number of individual animals from several  
different pig lines. Analyses of growth and performance test  
records showed significant associations of MC4R genotypes  
with backfat, growth rate and feed intake in a number of  
lines. It is probable that the variant amino acid residue of  
the MC4R mutation causes a significant change of the MC4R  
function. These results support the functional significance  
of a pig MC4R missense mutation and suggest that comparative  
genomics based on model species may be equally important for  
application to farm animals as they are for human medicine.

Identification of mutations in the *leptin* and the *leptin*  
*receptor* has provided some information on genetic components  
involved in the regulation of energy balance (Zhang et al.  
1994; Tartaglia et al. 1995). Genetic studies using animal  
models have facilitated the identification of major genetic  
causes of obesity (Andersson 1996; Pomp 1997; Giridharan  
1998). Furthermore, several other genes involved in the  
neural signaling pathway of energy homeostasis have been  
identified (Flier and Maratos-Flier 1998; Schwartz et al.  
1999). Of particular interest among candidate signaling  
molecules involved in the regulation of energy homeostasis is  
the melanocortin-4 receptor (MC4R). The MC4R response to



leptin signaling is a link between food intake and body weight (Seeley et al. 1997; Marsh et al. 1999). Neuropeptide Y (NPY) signaling in the central nervous system is also mediated by the MC4R protein (Kask et al. 1998). Several mutations in MC4R including frameshift and nonsense mutations are associated with dominantly inherited obesity in humans (Vaisse et al. 1998; Yeo et al. 1998). Some other MC4R missense mutations in humans have also been identified (Gotoda et al. 1997; Hinney et al. 1999) but the functional significance of these mutations has not been characterized.

Selection based on growth characteristics has been of great importance to the pig industry because of costs associated with feeding and consumer preference for lean meat. Efficient genetic improvement in these quantitative traits may be augmented through the use of marker assisted selection (MAS) using high density genetic maps (Dekkers and van Arendonk 1998; Rothschild and Plastow 1999). An important tool in this process is comparative mapping using the well-developed human and mouse gene maps, which assist in the identification of corresponding genomic regions or major genes controlling growth and performance traits in the pig. Biological understanding of complex traits in human or model species offers an alternative approach to identify genes responsible for the traits of economic interest in livestock. Several quantitative trait loci (QTL) linkage scans using phenotypically divergent breeds and candidate gene analyses have been successfully conducted for fatness and growth traits (Yu et al. 1995; Casas-Carrillo et al. 1997; Knorr et al. 1997; Knott et al. 1998; Rohrer et al. 1998; Wang et al. 1998; Paszek et al. 1999), but no individual genes with major effects on growth and performance traits have yet been established for commercial populations. The role of MC4R in feed intake and obesity suggests it may be an important genetic marker for the growth-related traits in the pig.

## Materials and Methods

Animals. Pigs were raised under normal production conditions under the care of PIC employees in nucleus farms in the United States and Europe.

PCR amplification of a pig MC4R gene fragment. Primers were designed from homologous regions of human and rat MC4R sequences (GenBank accession no. s77415 and u67863, respectively). The primers were: forward primer: 5'-TGG CAA TAG CCA AGA ACA AG-3' (SEQ. ID NO:6) and reverse primer: 5'-CAG GGG ATA GCA ACA GAT GA-3' (SEQ. ID NO:7). The PCR reaction was performed using 12.5 ng of porcine genomic DNA, 1x PCR buffer, 1.5 mM MgCl<sub>2</sub>, 0.125 mM dNTPs, 0.3 mM of each primer, and 0.35 U *Taq* DNA polymerase (Promega) in a 10μL final volume. The conditions for PCR were as follows: 2 min at 94°C; 35 cycles of 30 s at 94°C, 1 min at 56°C, 1 min 30 s at 92°C, and a final 15 min extension at 72°C in a Robocycler (Stratagene, La Jolla, CA).

Sequencing and mutation detection. Sequencing of the PCR products from several individual pigs of different breeds was conducted and the sequences were compared to detect any nucleotide change. Sequencing was performed on an ABI sequencer 377 (Applied Biosystems). The porcine MC4R sequence has been submitted to GenBank, and has accession number AF087937. The sequence analysis revealed one nucleotide substitution situated within a *TaqI* restriction enzyme recognition site (Kim et al. 1999). A set of primers was then designed to generate a smaller MC4R gene fragment, which contained only one informative *TaqI* restriction site to specify the polymorphic site and to facilitate the PCR-RFLP test. These primers were: forward 5'-TAC CCT GAC CAT CTT GAT TG-3' (SEQ. ID NO:10) and reverse: 5'-ATA GCA ACA GAT GAT CTC TTT G-3' (SEQ. ID NO:11).

## Results

Identification of a missense mutation in the pig MC4R gene. The MC4R gene consists of approximately 1 kb of coding sequence contained within a single exon. About 750 bp of a pig MC4R gene fragment was produced by PCR (Kim et al. 1999). The sequence of the PCR product confirmed that the PCR product is the MC4R gene with 92.2% and 97.6% identities at nucleotide and the amino acid levels, respectively, to the human MC4R sequence. Multiple alignments of the sequences from individual animals of several breeds identified a single nucleotide substitution (G→A; Fig. 5). The polymorphism revealed a missense mutation that replaces aspartic acid (GAU) with asparagine (AAU) at the position identical to amino acid 298 of human MC4R protein. To confirm this base change, we designed pig-specific primers flanking the polymorphic site and analyzed the polymorphism as a *TaqI* PCR-RFLP gel (Fig. 6). Figure 6 shows a *TaqI* digestion of the PCR product analyzed by agarose-gel electrophoresis. Allele 1 produced 156 and 70 bp fragments and allele 2 produced a 226 bp fragment as the PCR-RFLP. The heterozygote has both allele 1 and 2 fragments. Molecular marker (M) and MC4R genotypes are indicated at the top of each lane.

The MC4R missense mutation is within a highly conserved region among melanocortin receptors (MCR). The MCR is a subfamily of G-protein coupled receptors (GPCR) containing certain conserved structural elements common to most other GPCRs, but overall amino acid identities between MCR and other GPCRs are low (Tatro 1996). A multiple-alignment of the predicted amino acid sequences of the pig MC4R with MC4R proteins from other species, other MCR proteins, or representative GPCRs showed that the aspartic acid found at position 298 of the seventh transmembrane domain is very highly conserved in the MCR proteins (Fig. 7). It is interesting to note, however, that this position is occupied by asparagine in most other GPCRs. The MCR proteins show 40-80% amino acid identity with each other (Tatro 1996), but the

second intracytoplasmic loop and the seventh transmembrane domain are highly conserved among MCR proteins (Gantz et al. 1993). Some of the relationships between MCR structure and function have been discovered by the studies of natural and experimental mutations in humans and mice (Robbins et al. 1993; Valverde et al. 1995; Frandberg et al. 1998). These studies indicate that some mutations in highly conserved regions cause structural changes and alter the function of the receptor. The Asp298Asn substitution mutation could have an effect on the function of the receptor. However, this will require further testing but it is known that change of the homologous residue in MC1R (Asp294His) is associated with fair skin and red hair in humans (Valverde et al. 1995).

### EXAMPLE 3

Quantity and quality are descriptive terms of great importance in the meat industry. As the live animal is converted to meat and the meat moves along the line of distribution, from slaughters and processors to retailers and finally to consumers, the factor of quality becomes increasingly more important. Obviously, economic considerations influence the concerns for quantity and quality.

The condition of pale, soft and exudative (PSE) pork and generally very high variability of pork quality was recognized and documented by 1960, and both quality "defects" have been viewed as having less value for further processing and being inferior for consumers. Although an enormous amount of research has been directed at the problem through a half-century of effort, surveys of incidence showed, in pork produced in the U.S.A., that 18% was of inferior quality (PSE) in 1963 and 16% in 1992. Thus, the existence of gene markers associated with both the ability to change the levels of traits (i.g. meat color, water holding capacity, tenderness or marbling) as well as to reduce variation in

meat quality characteristics provides excellent opportunities for a dramatic improvement in meat quality. First, gene markers allow for significant steps to be made in the desired direction of quality traits; (e.g. improving technological yield of processed ham and reducing moisture (purge) losses of fresh ham and loins, by selecting against the RN gene in pigs). Secondly, gene markers will help reduce meat quality variation, since we can fix relevant genes in the breeding populations.

10 Meat quality is typically measured in slaughter plants in terms of the pH of the meat, color (using several different instruments and methods e.g. Minolta, [Min]), marbling and drip loss.

For example, the following descriptions of desirable meat quality characteristics are generally accepted by the industry based on their economic value at different segments of the pork supply chain:

5        Loin Minolta Lightness (L\*): The range of 43-47 units (from darker to lighter color) is acceptable, but L\* of 43 is better; i.e., has higher economic value, in general in this range\*\*

10       Loin Japanese Color Score (JCS): The range of 2.5 - 5.0 units (from lighter to darker color) is acceptable, but JCS of 3-4 is better

      Loin Marbling (level of intramuscular fat): Generally, higher marbling is better as it is associated with improved meat eating quality characteristics

15       Loin pHu: (ultimate meat acidity measured 24 hours post-mortem; this attribute is the single most important trait of pork quality); -- The range of 5.50 - 5.80 is desirable, but 5.80 is better as it positively influences the color and (low) purge of the meat

20       Ham Minolta lightness (L\*) The range of 43-52 units is acceptable, but lower (43) is better

      Ham pHu: higher; i.e., 5.80, is better

      Drip loss or purge: the range of 1%-3% is acceptable, but lower is better

25

\*\* this may be dependent upon market, for example in Japan darker pork is preferred. Sosnicki, A.A., E.R. Wilson, E.B. Sheiss, A. deVries, 1998 "Is there a cost effective way to produce high quality pork?", Reciprocal Meat Conference

30       Proceedings, Vol. 51.

## Results

Table 1

Least square means for different MC4R genotypic classes based on a sample of 1146 animals from six genetic lines (preferred class in bold)

Trait	Genotype			
	11	12	22	p value
Loin pHu	5.70	5.70	<b>5.73</b>	<0.01
Ham pHu	5.69	5.69	<b>5.72</b>	<0.07
Ham Min L	48.44	48.39	<b>47.38</b>	<0.03
Drip	2.29	2.43	<b>2.10</b>	<0.07
Loin Marbling	2.17	2.18	<b>2.25</b>	<0.42
Days to 110g*	169.2	168.5	<b>166.4</b>	<0.0001

\* The samples size for Days was 2366

Significant effects of marker genotype are identified for ultimate pH (pHu), color (Min) and drip loss and a desirable trend is observed for marbling. The size of the effects observed between genotypes while small are of commercial significance in terms of differences in meat quality. It can be seen from the results in Table 1 that allele 2 is the preferred allele in this sample for all four meat quality measures. Interestingly, this is the preferred allele for growth as reported in WO 00/06777. This is a particularly important finding, as it is somewhat unexpected. In general, there is a negative correlation between growth rate and meat quality. Indeed, there is a general perception that meat quality has decreased as breeders have selected for increased growth rate.

In some situations we might anticipate that the associations between the marker genotype and the traits may differ in direction. This will be the case where the marker utilized here is linked to the polymorphism or gene that is causing the effect. In this situation, the MC4R marker will still have utility, once the association has been identified by experimentation.

The traits measured here are only some of the measures that may be used for determining meat quality. Many others can be used that are correlated to these measures. Thus it will be expected that similar effects will be observed for such economically important traits as water holding and binding capacity, curing and cooking yields and that these traits and that these will also extend to related measures of eating quality such as tenderness, juiciness, flavor and taste. See Sosnicki, supra.

The present invention concerns the identification of significant associations between the MC4R marker genotype and meat quality. It will be realized by those skilled in the art that other gene markers located in this region of the swine genome (swine chromosome 1) will also be suitable for marker assisted selection of these traits.

#### EXAMPLE 4

A total of 257 animals from a Pietrain-based line of pigs were slaughtered and meat quality characteristics determined at the time of slaughter and during post-slaughter handling/conditioning for meat production. MC4R genotypes were determined using methods disclosed herein. Associations between marker genotype and MQ traits were then calculated. The results are depicted in Table 2.

Significant associations were obtained for ultimate pH and color (Minolta L) of the ham. Hams from animals of genotype 2,2 will be preferred in markets which favor these characteristics.

The association can be used to select the parents of slaughter pigs or to improve breeding stock by within line selection. Alternatively, ham processors may chose to purchase pigs of the preferred genotype in order to improve the overall quality of the product (2,2 carcasses will be a better color and will be expected to provide greater yield than those from 1,1 or 1,2 animals). In addition, by selecting a single genotype they will also reduce the



variation in product quality due to the different MC4R genotypes.

5

Table 2

MC4R genotype and Ham pHu and Min L for a Pietrain based line selected to be free of the Halothane gene

Genotype	n	pHu	Min L
11	119	5.72	49.70
12	101	5.73	50.03
22	37	<b>5.80</b>	<b>47.83</b>
p		<0.04	<0.09

10

The present invention describes an association between MC4R genotype and meat quality characteristics such as pH, color and marbling in the pig. These traits are in turn associated with visual appearance and processing and eating quality characteristics such as tenderness.

15

These traits also describe meat quality in other species such as beef and lamb. Because of the relatively close evolutionary link between pigs and other meat species it can be predicted that variation in this gene is also likely to be associated with meat quality (MQ) in these other species. Polymorphisms can be identified in the MC4R gene of these species using the same approach set out here and the resulting SNPs used for association analysis.

20

Having described the invention with reference to particular compositions, theories of effectiveness, and the like, it will be apparent to those of skill in the art that it is not intended that the invention be limited by such illustrative embodiments or mechanisms, and that modifications can be made without departing from the scope or spirit of the invention, as defined by the appended claims. It is intended that all such obvious modifications and variations be included within the scope of the present invention as defined in the appended claims. The claims are

25

30

meant to cover the claimed components and steps in any sequence which is effective to meet the objectives there intended, unless the context specifically indicates to the contrary.

The following citations are hereby incorporated in their entirety by reference:

- Andersson LB (1996), Ann Med., 28:5-7.
- Arden KC (1990), Cytogenet. Cell Genet. 53:161-165.
- 5 Barinaga M (1996), Science 271:913.
- Bray GA (1978). Physiol. Rev. 59, 719-809.
- Casas-Carillo E (1997), J Anim Sci 75:2047-2053.
- Chajlani V, Biochem. Biophys. Res. Commun., 195, 866-873.
- Chen H (1995), Cell 84, 491-495.
- 10 Chua SC (1996). Science 271, 994-996.
- Chung WK (1996). Genome Res. 6, 431-438.
- Cioffi JA (1996). Nature Med. 2, 585-589.
- Cybulsky MI (1991), Proc. Nat. Acad. Sci. 88:7859-7863.
- 15 Dekkers JCM (1998), Genet Res 71:257-275.
- Flier JS (1998), Cell 92:437-440.
- Frandberg P (1997), Biochem Biophys Res Commun 236:489-492.
- Frankel WN (1996), Nat Genet 14:371-373.
- Gantz (1993), J. Biol. Chem. 268, 15174-15179.
- 20 Gantz (1994), Biochem. Biophys. Res. Comm., 200, 1214-1220.
- Gantz I (1993a), J. Biol. Chem. 268, 8246-8250.
- Giridharan NV (1998), Indian J Med res 108:225-242.
- Gotoda T (1997), Diabetologia 40:976-979.
- Goureau A (1996). Genomics.36:252-262.
- 25 Griffon N (1994), Biochem. Biophys. Res. Comm., 200, 1007-1014.
- Helm J (1994), J. Anim. Science 72:2764.
- Hinney A (1999), J Clin Endocrinol Metab 84:1483-1486.
- Huszar (1997), Cell 88:131-141.
- 30 Kask A (1998), Biochem Biophys Res Commun 248:245-249.
- Kim KS (2000), J Anim Sci 78:791-792.
- Knorr C (1997), Anim Genet 28:124-128.
- Knott SA (1998), Genetics 149:1069-1680.
- Labbe O (1994), Biochemistry, 33, 4543-4549.
- 35 Lee GH (1996), Nature 379, 632-635.
- Lu Dongsi (1994), Nature:371:799-802
- Marsh DJ (1999), Nat Genet 21:119-122.

- Meuwissen, T.H.E. and Goddard, M.E. (1996) "The use of Marker Haplotypes in Animal Breeding Schemes", Genet. Sel. Evol., 28 161-176
- 5 Montjoy KG (1992), Science 257, 1248-1251.  
 Montjoy KG (1994), Mol. Endocrinol. 8, 1298-1308.  
 Paszek A (1996), Proceed. Midwest. ASAS Meeting, p. 24.  
 Paszek AA (1999), Mamm Genome 10:117-122.  
 Phillips M (1996), Nature Genetics 13:18-19.
- 10 Pomp D (1997), Behav Genet 27:285-306.  
 Robbins LS (1993), Cell 72:827-834.  
 Robic A (1996), Mamm. Genome 7, 438-445.  
 Rohrer GA (1998), J Anim Sci 76:2247-2254.  
 Rothschild MF (1996), PNAS 93: 201-205.
- 15 Rothschild MF (1999), AgBiotechNet 1:1-8.  
 Schwartz MW (1999), Am J Clin Nutr 69:584-96.  
 Seeley RJ (1997), Nature 390:349.  
 Tartaglia LA (1995), Cell 83:1263-1271.  
 Tatro JB (1996), Neuroimmunomodulation 3:259-284.
- 20 Truett GE (1995), Mamm. Genome 6, 25-30.  
 Vaisse C (1998), Nat Genet 20:113-114.  
 Valverde P (1995), Nat Genet 11:328-330.  
 Wang L (1998), J Anim Sci 76:2560-2567.  
 Warner CM (1991), Immunogenetics of the MHC, VCH Publishers, NY, NY, pp. 368-397.
- 25 Winick JD (1996), Genomics 36, 221-222.  
 Xy Y (1994), Biol. Reprod. 51:695-699.  
 Yeo GS (1998), Nat Genet 20:111-112.  
 Yerle M (1996), Cytogenet. Cell Genet. 73, 194-202.
- 30 Youngs CR (1993), J. Anim. Sci., 1561-1565.  
 Yu TP (1995), J Anim Sci 73:1282-1288.  
 Zhang Y (1994), Nature 372:425-432.  
 Ziang ZH (1999), Mamm Genome 12:191-193.